


Maths Learning Service: Revision **Mathematics IA**  
**Complex Numbers**



The *imaginary number*  $i = \sqrt{-1}$  is an extension to the real number system which allows us to solve equations such as

$$x^2 = -1.$$

A *complex number* is any number of the form  $z = a + bi$ , where  $a$  and  $b$  are real numbers.

**Note:** All numbers involving  $i$  can be written in this form.

**Examples:** (a)  $i^2 + i^3$  (b)  $\frac{2i - 3}{i + 1}$

$$= -1 + i^2 i = -1 - i$$

$$= \frac{2i - 3}{i + 1} \times \frac{i - 1}{i - 1}$$

$$= \frac{2i^2 - 5i + 3}{i^2 - 1}$$

$$= \frac{-2 - 5i + 3}{-1 - 1}$$

$$= \frac{1 - 5i}{-2} = -\frac{1}{2} + \frac{5}{2}i$$

- Notes:** (1) In  $z = a + bi$ ,  $a$  is the *real part* of  $z$ .  
 $b$  is the *imaginary part* of  $z$ .  
 (2) If  $b = 0$ ,  $z$  is a real number.  
 If  $a = 0$ ,  $z$  is a *purely imaginary number*.

$z = a - bi$  is the *complex conjugate* of  $z = a + bi$ . The solutions to quadratic equations are complex conjugates.

**Example:** Solve  $x^2 - 2x + 10 = 0$ .

**Solution:** Using the quadratic formula with  $a = 1$ ,  $b = -2$  and  $c = 10$ ,

$$x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4 \times 1 \times 10}}{2 \times 1}$$

$$= \frac{2 \pm \sqrt{-36}}{2}$$

$$= \frac{2 \pm \sqrt{36 \times -1}}{2}$$

$$= \frac{2 \pm 6i}{2}$$

$$= 1 - 3i \text{ or } 1 + 3i.$$


## Exercises

- (1) Rewrite the following in the form  $a + bi$ :
- (a)  $(2i + 6) + (5i - 1)$       (b)  $(3i + 2)(i - 1)$
- (c)  $\frac{i}{1 + 2i}$       (d)  $\frac{1}{2 - i} + \frac{2}{2 + i}$
- (2) If  $z_1 = 1 - 2i$  and  $z_2 = 2 + i$ , find:
- (a)  $z_1 + z_2$       (b)  $\overline{z_1 + z_2}$
- (3) Solve for  $x$ :
- (a)  $x^2 - 10x + 29 = 0$       (b)  $\frac{1}{x} = 1 - 2x$

## The Complex (Argand) Plane

The complex number  $z = a + bi$  can be represented on a number *plane* (rather than a number *line*) with co-ordinates  $(a, b)$ . The  $x$ -axis represents the real component of  $z$  and the  $y$ -axis represents the imaginary component.

Examples: (a)  $3 + 2i$       (b)  $3 - 2i$       (c)  $-2$       (d)  $4i$



The distance of a complex number from the origin of the Argand Plane is called the *modulus* of the complex number  $z$  (or  $|z|$ ). By Pythagoras' Theorem:


$$|z| = \sqrt{a^2 + b^2}.$$

Examples: (a)  $|3 + 2i|$       (b)  $|3 - 2i|$       (c)  $|-2|$       (d)  $|4i|$

$$= \sqrt{3^2 + 2^2} \qquad = \sqrt{3^2 + (-2)^2} \qquad = \sqrt{(-2)^2 + 0} \qquad = \sqrt{0 + 4^2}$$

$$= \sqrt{13} \qquad = \sqrt{13} \qquad = 2 \qquad = 4$$





(The reader is left to investigate why this solution works geometrically.)

**Exercises**

(4) Find:

- (a)  $|1 + i|$                       (b)  $|1 - i|$                       (c)  $|-6i|$

(5) Find all complex numbers  $z = x + iy$  which satisfy:

- (a)  $|z + 1| = 1$                       (b)  $|z| = |z + 1|$

**Answers to Exercises**

- (1) (a)  $5 + 7i$                       (b)  $-5 - i$   
 (c)  $\frac{2}{5} + \frac{1}{5}i$                       (d)  $\frac{6}{5} - \frac{1}{5}i$

- (2) (a)  $3 + i$                       (b)  $3 + i$   
 (This example demonstrates the property that  $\overline{z_1 + z_2} = \overline{z_1} + \overline{z_2}$ .)

- (3) (a)  $5 - 2i$  or  $5 + 2i$                       (b)  $\frac{1}{4} - \frac{\sqrt{7}}{4}i$  or  $\frac{1}{4} + \frac{\sqrt{7}}{4}i$

- (4) (a)  $\sqrt{2}$                       (b)  $\sqrt{2}$                       (c) 6

- (5) (a)  $(x + 1)^2 + y^2 = 1$                       (b)  $x = -\frac{1}{2}$   
 (A circle of radius 1 centred on  $(-1, 0)$ )                      (A vertical line)

