

# DRY DEPOSITION OF POLLUTANT AND MARKER PARTICLES ONTO LIVE MOUSE AIRWAY SURFACES REVEALS HETEROGENEOUS MUCOCILIARY TRANSIT BEHAVIOUR

Martin Donnelley<sup>1,2,3,4</sup>, Karen Staley<sup>1,2,3,4</sup>, Kaye Morgan<sup>2</sup>, Andreas Fouras<sup>5</sup>, David Parsons<sup>1,2,3,4,6</sup>

1. Respiratory and Sleep Medicine, Women's and Children's Hospital, South Australia
2. Women's and Children's Health Research Institute, South Australia
3. School of Paediatrics and Reproductive Health, and 4. Centre for Stem Cell Research, University of Adelaide, South Australia
5. School of Physics, 6. Monash Biomedical Imaging, and 7. Division of Biological Engineering, Monash University, Victoria
8. Australian Synchrotron, Victoria



Government of South Australia  
SA Health



## BACKGROUND:


- Airborne pollutants are a considerable health concern and have the potential to impact on respiratory diseases such as cystic fibrosis (CF)
- We have previously described synchrotron imaging techniques for examining the mucociliary transit (MCT) behavior of particulates in the nasal airways and trachea of anaesthetised mice
- Our previous studies delivered particulates in a fluid bolus, however the presence of the carrier fluid perturbed the airway surface and obscured the manner of deposition
- The aim of this study was to verify that synchrotron phase contrast X-ray imaging (PCXI) can be used to detect, monitor and compare the deposition and MCT behaviour of pollutant and marker particles after dry deposition into the trachea of live mice

## METHODS:

- Intubated (flexiVent ventilation) C57Bl/6 mice (n=8) were imaged on the BL20XU beamline at the Spring-8 synchrotron in Japan
- Particles of borosilicate glass, quarry dust and lead ore, as well as reference 14 µm hollow glass beads were deposited into the trachea using a nebuliser pump (PennCentury, Wyndmoor, PA, USA) at a rate of one per breath (in an end-inspiratory pause) for five minutes using a high-resolution camera
- Images were enhanced post experiment (Matlab, The Mathworks, MA, USA) assembled into movies showing particle movement and analysed using ImageJ (National Institutes of Health, Bethesda, MD, USA)

## RESULTS:

- The first movement of particulates tended to appear approximately 2-3 minutes after delivery
- As in previous studies the particle transit was heterogeneous: after deposition some particles did not move, while others transited the field of view rapidly
- The big and heavy particles lead and tantalum moved substantially less than the other particulates



- Most particles did not follow a linear path along the airway surface, many followed seemingly random paths
- There was more movement of particles along the bottom quarter of the tracheal surface than the remainder of the trachea, possibly due to the quantity of mucus present and gravitational effects
- In some cases contraction of the tracheal airway throughout the imaging period

FIG 1. Particle visibility on the airway surface of four mice (a) silver coated hollow glass beads, (b) lead, (c) fibreglass fibres, and (d) quarry dust. In these static images it is hard to clearly identify some of the particles (excluding the large lead particles and fibreglass fibres).

the imaging location is just above the main carina.

FIG 2. Particle movement vectors from two animals (a) and (b) are shown in the sites of all lead particles and (d) on the airway surface over the 5 minute imaging period. The red and white regions contained the most particle movement. Movement in the top third of the frame

Figure (c) were the only particles that moved. On average, there is more particle MCT along and towards the dorsal surface. A grid of the movement vectors (green arrows) is overlaid on the image. Engineering flow measurement technique – streamlines.

Figure orientation and location is the same as Figure 1.

## CONCLUSION:

Synchrotron PCXI permits detection of particle transit via MCT along live mouse trachea. We are continuing with studies to improve our direct and non-invasive MCT assessment methods to assist our understanding and treatment of respiratory diseases such as CF.

## ACKNOWLEDGEMENTS:

Supported by NH&MRC Australia and the U.S. National Science Foundation. Spring-8 experiments performed under proposal 2010B1137. Travel funding provided by the Australian Synchrotron ISA Program. We thank Naoto Yagi and Kentaro Uesugi for their assistance.